

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/276303819>

THE OCCURRENCE PROBABILITIES OF ROGUE WAVES IN DIFFERENT NONLINEAR STAGES

Article in Coastal Engineering Proceedings · January 2014

DOI: 10.9753/icce.v34.waves.35

CITATIONS

0

READS

24

5 authors, including:



Aifeng Tao

Hohai University

12 PUBLICATIONS 28 CITATIONS

SEE PROFILE

All content following this page was uploaded by [Aifeng Tao](#) on 29 February 2016.

The user has requested enhancement of the downloaded file. All in-text references [underlined in blue](#) are added to the original document and are linked to publications on ResearchGate, letting you access and read them immediately.

THE OCCURRENCE PROBABILITIES OF ROGUE WAVES IN DIFFERENT NONLINEAR STAGES

Aifeng Tao^{1,2}, Keren Qi^{1,2}, Jinhai Zheng^{1,2}, Ji Peng^{1,2}, Yuqing Wu^{1,2}

The occurrence probabilities of Rogue Waves in different nonlinear states are investigated based on high-order spectral method (HOSM), which is a direct phase-resolved numerical method. The focus is given to the occurrence probability of Rogue Waves in the nonlinear evolution stage where the Benjamin-Feir Instability is not the dominant mechanism due to the quartet resonance interactions for wave fields with a broad range in frequencies. The initial wave trains are generated from Stokes waves and two sidebands. Based on the simulation, we find that the Kurtosis evolves distinctly at three nonlinear stages and shows a weak relation to the probabilities of the Rogue Waves. We also introduce a simple Entropy formula, turning out to close a stable value.

Keywords: Rogue Waves, HOS, Nonlinear mechanism, probability, Kurtosis, Entropy

1. Introduction

The concept of Rogue Wave was first put forward by Draper (1965) to describe exceptionally high waves surprisingly appearing on the sea surface, which may cause disastrous damages to offshore structures and surface vessels. One typical and common operational definition of Rogue Wave defines that its wave height should exceed the significant wave height in 2 times. In the recent two decades, a large number of Rogue Wave studies have been conducted theoretically, numerically, experimentally and based on field data, which have significantly advanced our knowledge of this kind special and notorious disastrous wave. Some new results including the discussion and debate for the Rogue Waves have been summarized by [Akhmediev & Pelinovsky \(2010\)](#) in the special topics of the European Physical Journal. Due to the potential risk, the Rogue Waves have attracted the attention of the shipping and offshore industry. So that some big international research projects have been initiated as mentioned by [Bithner-Gregersen & Toffoli \(2012\)](#). However, there is still not a unifying concept towards the Rogue Waves can be concluded. The occurrence probability of Rogue Waves is one of the most interesting topics since it related to the engineering design practice directly. In the last decades, a number of works have been done in this field, such as Mori (2006), Akhmediev (2011), Tomas (2012), Xiao (2013), Wang(2014) based on milestone work conducted by Janssen (2003), where the Benjamin-Feir-Index(BFI) was introduced and explained theoretically. However, physically, BFI is useful because it is a parameter showing the conditions of the Benjamin-Feir Instability. If a wave train can propagate in a large scale distance, or a sea state can evolve for a long time, such as the ocean in the monsoon seasons, the nonlinearity may grow to a stage beyond the Benjamin-Feir Instability can control, since the quartet resonance interactions for wave fields with a broad range in frequencies and directions. The effectiveness of BFI characterizing the occurrence probability of Rogue Waves seems to be a problem.

One of The latest studies for the probability of Rogue Waves has been performed by [Bithner-Gregersen & Toffoli \(2012\)](#), focusing on the design practice. One of their results is that ‘the extreme crest considerably increases for prolonged sea state durations’. Considering the concept, which indicates that there are different kinds of Rogue Waves according to their intrinsic nonlinear mechanisms, shown by Liu (2006) based on field data and [Tao\(2011, 2012a, 2012b\)](#) via numerical simulations. It does need carry out some studies to understand the characteristics of the Rogue Waves in a long time evolved wave state, including the occurrence of them.

2. Approach

After a systemically verification with experimental data provided by [Shemer & Sergeeva\(2009\)](#), the high-order spectral method(HOSM) developed by [Dommermuth & Yue\(1987\)](#) is applied for direct phase-resolved simulation of nonlinear random wave fields evolution. HOSM resolves the phase of a large number (N) of wave modes and accounts for their nonlinear interactions up to an arbitrary high order (M) including broadband non-resonant and resonant interactions up to any specified order. Meanwhile viscous dissipation and wave breaking dissipation are modelled in domain. Due to its high efficiency and accuracy, the HOSM is an effective approach for long-time and large-space simulation of nonlinear wave-field evolutions.

3. Modulated Stokes wave train long time evolution

In this section, we will investigate the characteristics of long time evolution processes generated from modulated Stokes wave train. First we studied the evolution process of narrow band modulated wave train, constructed by a Stokes carrier wave and imposed sideband according to most unstable modulation instability condition. Then the characteristics of extreme waves will be discussed to introduce the evolutions of different parameters.

¹State Key Laboratory of Hydrology-Water Resources and Hydraulic Engineering, Hohai University, Nanjing 210098, China

² College of Harbor Coastal and Offshore Engineering, Hohai University, Nanjing 210098, China

3.1 Initial condition with weakly modulated wave train

For the narrow band modulated wave train, the initial conditions we used are as following.

$$\left. \begin{aligned} \eta(x,0) &= \eta[\varepsilon_0, k_0] + r_1 a_0 \cos(k_- x - \theta_-) + r_2 a_0 \cos(k_+ x - \theta_+) \\ \phi^s(x,0) &= -\phi^s[\varepsilon_0, k_0] + \frac{r_1 a_0}{\sqrt{k_-}} e^{k_- \eta} \sin(k_- x - \theta_-) + \frac{r_2 a_0}{\sqrt{k_+}} e^{k_+ \eta} \sin(k_+ x - \theta_+) \end{aligned} \right\}$$

where $\eta(x,0)$ and $\phi^s(x,0)$ are respectively the free-surface elevation and potential of a right-going Stokes wave of steepness ε_0 and wave number k_0 . Here $\varepsilon_0 = k_0 a_0$ and a_0 is the surface elevation of carrier wave. $k_{\pm} = k_0 \pm \Delta k$ and θ_{\pm} are wave numbers and phases of sideband respectively. Those parameters are chosen as the most unstable conditions in the initial period. The initial wave relative amplitude spectrum and wave surface for $\varepsilon_0 = 0.05$ are shown in Fig.1. For all the cases, the calculation domain is $200l_0$. The parameters for HOS are $M=7$, $N=8192$, $T_0/dt = 128$. Here dt is the time step for simulation.

Cases	M	$\Delta k / k_0$	θ_{\pm}	$r_{1,2}$	Evolution time (t/T0)
I	7	0.1	$\pi / 4$	0.1	30000




Fig.1. Initial wave relative amplitude spectrum (left) and wave surface (right) for $\varepsilon_0 = 0.05$

Using HOS method, calculation accuracy would significantly enhances as wave modes (N) being incredibly close to nonlinear wave order (M). Higher the nonlinear wave order (M), smaller the round-off errors. However, higher precision requiring higher demand of computing makes it necessary for us to determine the suitable M for this research. The variable E/E_0 shown in Fig.2 indicates the conservation of the wave train with minute energy dissipation (only less than 0.01%) over evolution time. As shown in the picture, it has smooth evolution process, assuring the numerical results.




Fig.2 Energy evolution of Stokes wave train for $\varepsilon_0 = 0.05$, $r_1=0.1$

3.2 Characteristics of Freak waves and the related mechanisms

3.2.1 Kurtosis

Characteristic parameters such as Kurtosis, occurrence probability, etc. change as waves propagate and the Kurtosis is calculated as

$$Kurtosis(t_i) \equiv \frac{1}{N_x} \sum_{j,t=t_i} \eta_j^4 / \eta_{rms}^4, \quad \eta_{rms} = \sqrt{\frac{1}{N_x} \sum_i \eta_i^2}$$

where η is the sea level displacement with zero mean level, η_{rms} is root-mean-square value of η , and N_x is the number of values. Kurtosis describes the degree that actual probability distribution deviates from Gaussian distribution with the value of 3. In this paper, the Kurtosis is calculated based on the wave surface elevation of whole calculation domain for each carrier wave period T_0 . From Fig.3, it can be seen clearly that the Kurtosis presents distinct different properties in different nonlinear stages.

The Kurtosis tends to increase gradually and the average value is greater than 3 beyond the B-F instability dominated stage.




Fig.3.The evolution of Kurtosis and the average trend

Fig.4 is given to find out the relationships between Kurtosis and extreme waves in each time scale. Interestingly, there appears to have distinct trends at different stages of evolution. In Fig.5, the relationship between the Kurtosis and the extreme waves at the first stage ($0 < t/T_0 \leq 7100$) can be given as a fitting equation ($y = 0.5052e^{0.9582x}$ with correlation coefficient of $R=0.9875$) which is close to exponential manner while the relationship at the third stage ($9000 < t/T_0 \leq 30000$) shows a linear trend with much lower correlation. There is also an intermediate stage ($7100 < t/T_0 \leq 9000$) between the two physical nonlinear ones. It can be deduced that the Kurtosis is close related to the extreme waves in different nonlinear stages.




Fig.4.Relationship between the Kurtosis and the extreme waves




Fig.5. Relationships between the Kurtosis and the extreme waves at three nonlinear stages

3.2.2 Occurrence probability

Occurrence probabilities of the Rouge Waves, which are calculated based on the wave surface elevation of whole calculation domain at each period, evolve with time as left picture of Fig.6 shows. It could be perceived that the probabilities of Rouge waves are always large at the peak of modulation. In order to catch the trend, the probabilities are averaged for each 3000 T_0 shown in the right picture, which become stable and present increase trend beyond the B-F instability dominated stage.




Fig.6. The evolution of occurrence probabilities of Rouge Waves and its average trend

We try to find parameters which related to the probability, the results are shown in Fig.7. The relationship between probability of Rouge Waves and extreme wave heights is obscure, which means higher Rouge Waves and high probabilities may not occur together, while the probabilities weakly related to the Kurtosis, which means the probabilities have a trend to increase with high nonlinearity.




Fig.7. Relationships between occurrence probability and other parameters (extreme waves and the Kurtosis respectively, from left to right)

To further study the varying pattern of probabilities of Rouge Waves based on fix point wave surface, we set four fix points working as anchored buoys (shown in Fig.8). We calculated the probabilities based on the wave surface time series on spot1 for each 128 T_0 and the average values are calculated from each 16 probabilities. Clearly to see from Fig.8, the probabilities from fix points present a similar trend with those from whole calculation domain.




Fig.8.Fix points setting schematic and the evolution of probabilities based on fix point wave surface (from left to right)

3.2.3 Entropy

After initial wave train evolves over enough long periods, it processes into random waves. In order to investigate the fundamental changes for the whole wave field, a simple Entropy formula is introduced to express the generated random waves. Here, Entropy is calculated as

$$E = \frac{-\sum p \ln(p)}{\ln(N)}$$

Wave heights are counted applying the zero-up (or zero-down) crossing method and divided into several intervals. In the expression above, p presents the probability that wave heights lie in a certain interval and N is the number of intervals. Entropy ranges from 0 to 1 while the value corresponding to the actual ocean waves falls in between. The Entropy of Gauss spectrum and JONSWAP spectrum are 0.29 and 0.39 respectively. The Entropy of our case is increasing along with the evolution process from imposed wave train to random waves as shows in Fig.9. Its tendency, being to close a stable value, may relate a particular wave spectrum.




Fig.9.Evolution of the Entropy of wave heights

4 Conclusion

We consider a very long-time evolution of modulated Stokes wave trains, which is obtained using nonlinear HOS simulation (with order $M=7$). On the basis of simulated results, the Kurtosis is close related with extreme waves in the whole long time evolution processes while the relationships are different distinctly for three nonlinear stages. For the probabilities of Rouge Waves, it presents a weak trend to increase with evolution time going on. Finally, by introducing a simple Entropy, we find the wave field displays a trend to be random waves and the wave spectrums tend to be a stable formula.

Acknowledgments

This research work is funded by the National Natural Science Fund (41106001, 51137002), the National Science Fund for Distinguished Young Scholars (51425930), the National Key Technology Research and Development Program (2012BAB03B01), the 111 Project (B12032), the Qing Lan Project of Jiangsu Province, the 333 Project of Jiangsu Province

(BRA2012130), the Scientific Research Fund for the Returned Overseas Chinese Scholars of the State Education Ministry ([2012]1707), the Natural Science Foundation Project of Jiangsu Province (BK2011026) and the Basic Research Fund From State Key Laboratory of Hydrology-Water Resources and Hydraulic Engineering, Hohai University (20145027512 and 20145028412).

REFERENCES

- A. S. Lev Shemer, (2009). "An experimental study of spatial evolution of statistical parameters in a unidirectional narrow-banded random wavefield." *Journal of Geophysical Research* 114(C01015): 1-11.
- A. Tao, J. Zheng, B. Chen, H. Li, J. Peng (2012). Propeties of Freak Waves induced by two kinds of nonlinear mechanisms. 33rd Conference on Coastal Engineering, Santander, Spain.
- A. Tao, J. Zheng, S. Mee Mee, B.Chen (2011). "Re-study on recurrence period of Stokes wave train with High Order Spectral method", *China Ocean Eng.*, 25, 679–686, 2011.
- A. Tao, J.Zheng, S. Mee Mee, B.Chen (2012). "The Most Unstable Conditions of Modulation Instability", *J. Appl. Math.*, 656873, 11pp., doi:10.1155/2012/656873, 2012a.
- A. Tomas, M. M., F.J. Mendez, G. Coco, and I.J. Losada (2012). "Predicting the ocurrence probability of freak waves baed on buoy data and non-stationary extreme value models." *Geophysical Research Abstracts* 14: 1.
- D. G. Dommermuth, and D. K. P. Yue (1987). "A high-order spectral method for the study of nonlinear gravity waves." *Journal of Fluid Mechanics* 184(1): 267-288.
- E. M. Bitner-Gregersen, and A. Toffoli (2012). "On the probability of occurrence of rogue waves." *Natural Hazards and Earth System Science* 12(3): 751-762.
- N. Akhmediev, A. Ankiewicz, J.M. Soto-Crespob, J.M. Dudleyc(2011). "Rogue wave early warning through spectral measurements?" *Physics Letters A* 375(3): 541-544.
- N. Akhmediev, and E. Pelinovsky (2010). "Editorial – Introductory remarks on “Discussion & Debate: Rogue Waves – Towards a Unifying Concept?”." *The European Physical Journal Special Topics* 185(1): 1-4.
- N. Mori, P. A. E. M. Janssen. (2006). "On Kurtosis and Occurrence Probability of Freak Waves." *J. Phys. Oceanogr.* 36: 1471-1483.
- P. A. E. M. Janssen, (2003). "Nonlinear four-wave interaction and freak waves." *J. Phys. Oceanogr.* 33: 2001–2018.
- Paul C. Liu, Keith R. MacHutchon (2006). Are there different kinds of Rogue Waves? Proceeding of the 25th International Conference on Offshore Mechanics and Arctic Engineering, Hamburg, Germany.
- Xiao, W., Y. Liu, G. Wual and Dick K. P. Yue (2013). "Rogue wave occurrence and dynamics by direct simulations of nonlinear wave-field evolution." *Journal of Fluid Mechanics* 720: 357-392.
- Y. Wang, A. Tao, J. Zheng, D. Doong, J. Fan, J. Peng (2014). "A preliminary investigation of rogue waves off the Jiangsu coast, China", *Nat. Hazards Earth Syst. Sci.*, 14, 2521-2527, doi:10.5194/nhess-14-2521-2014.